

UMDHEP 93-199

May 1993

Effect of CP-Odd Planck Scale Induced Operators on the Majoron Scale [†]

R.N. Mohapatra and X. Zhang

Department of Physics and Astronomy

University of Maryland

College Park, MD 20742

Abstract

We extend a recent work on the effect of non-perturbative Planck scale effects on models with spontaneously broken global B-L symmetry, where it was shown that scale V_{BL} of this symmetry must be less than 10 TeV. We show that, if the Planck scale effects break CP invariance, an allowed window for V_{BL} appears where $10^8 \text{GeV} \leq V_{BL} \leq 10^{11} \text{GeV}$.

[†] Work supported by a grant from the National Science Foundation

In recent months, there has been growing interest in the possibility[1] that non-perturbative gravitational effects may not respect any global symmetries of nature. Early work on “no hair” theorem[2] and recent analyses of the effect of “worm-hole” solutions on low energy particle physics seem to provide evidence for this speculation. In this brief report, we comment on the impact of this speculation on one particular class of extensions of the standard model where small neutrino mass arises in a natural manner, the singlet majoron model[3].

In a recent analysis[4] of the Planck scale effects on the singlet Majoron model, it was concluded that, there must be an upper bound on the V_{BL} , the scale of B-L symmetry breaking *i.e.*

$$V_{BL} < \left(\frac{m_\nu}{25\text{eV}}\right)^{4/7} \times 10 \text{ TeV}. \quad (1)$$

The basic idea is that, the Planck scale effects are expected in general to induce non-renormalizable terms of the following type in the low energy effective lagrangian:

$$\begin{aligned} V_{Planck} = & \frac{1}{M_{Planck}} (\lambda_1 \sigma^5 + \lambda_2 \sigma^4 \sigma^* + \lambda_3 \sigma^3 \sigma^{*2}) \\ & + \frac{1}{M_{Planck}} (\lambda_4 \phi^+ \phi \sigma^3 + \lambda_5 \phi^+ \phi \sigma^2 \sigma^*) \\ & + \dots \end{aligned} \quad (2)$$

where ϕ is the standard model higgs doublet, σ siglet field with B-L number -2. We parametrize $\sigma = \frac{1}{\sqrt{2}}(\eta + V_{BL})e^{i\chi/V_{BL}}$. This generates a non-zero

mass for the Majoron (χ):

$$m_\chi \simeq \left(\frac{25}{2}\lambda_1 + \frac{9}{2}\lambda_2 + \frac{1}{2}\lambda_3 \right)^{1/2} \left(\frac{V_{BL}}{V_{WK}} \right)^{3/2} \text{ keV}, \quad (3)$$

for $V_{BL} > V_{WK}$. If the Majoron is stable, clearly $V_{BL} > V_{WK}$ will be in contradiction with the cosmological mass density constraints. The situation does not improve, for unstable Majoron, because the only decay mode for the Majoron is to two neutrinos. The strength of this coupling is fixed by low energy considerations such as neutrino mass and is therefore quite constrained. More importantly, it was concluded in Ref.[4] that, the upper limit on V_{BL} is given by eq.(1). A slight refinement of this limit has recently been provided in Ref.[5].

In this report, we look at the impact of non-renormalizable operators on the Majoron decay. It is clear that for $V_{BL} \geq 10^8 \text{ GeV}$, eq.(3) implies that, $m_\chi \geq 1 \text{ TeV}$ implying that, it could perhaps decay to a pair of Higgs bosons (H). However, since the Higgs boson is CP-even and the Majoron is a CP-odd particle, $\chi \rightarrow n H$ is forbidden in the limit of CP- conservation. CP-violation is therefore necessary for any $\chi \rightarrow n H$ decay to occur. Furthermore, no such decay can occur from the renormalizable sector of the model. Therefore, we must look at CP-violating non-renormalizable Planck-induced operators. Two such operators are in eq.(2) if λ_4 and λ_5 are complex. They induce coupling of type:

$$\mathcal{L}_\chi = \sin \delta \ |\lambda_5| \ \frac{V_{BL}^2}{2M_{Planck}} \ \chi \ H^2 \ , \quad (4)$$

where we redefine the σ -field to absorb the CP-phase of λ_4 ; δ is the phase of λ_5 in this new basis. We limit the value of $|\lambda_5| \leq M_{Planck} V_{WK}^2 / V_{BL}^3$ in

order that the effective Higgs mass remains of the electroweak scale. Using this and $m_\chi \gg 2 m_H$, we get for the lifetime τ_χ of the Majoron:

$$\tau_\chi \simeq \frac{10^{-14}}{\sin^2 \delta} \left(\frac{V_{BL}}{10^8 \text{GeV}} \right)^{7/2} \text{sec}. \quad (5)$$

In order to see what constraints it implies on V_{BL} , let us look at the picture of the early universe with massive Majoron with $m_\chi \geq \text{TeV}$, which corresponds to $V_{BL} \geq 10^8 \text{ GeV}$. As was observed in Ref.[4], below $T = m_{\nu_R} \simeq V_{BL}$, the Majoron decouples and dilutes like a non-interacting gas. If its lifetime is longer than 10^{-1} sec or so, it will contribute an enormous amount to the energy density at the epoch of nucleosynthesis *i.e.*

$$\rho_\chi(T = 1\text{MeV}) \simeq Y n_\gamma m_\chi, \quad (6)$$

where the factor Y is the dilution factor for the Majoron density and $Y \simeq \frac{1}{10}$ [6]. This will significantly alter the predictions for Helium abundance.

On the other hand, if χ decays before $t_U = 10^{-1} - 10^{-2} \text{sec}$. the decay products $\chi \rightarrow 2 H \rightarrow 2b \ 2\bar{b} \rightarrow 2c \ 2\bar{c} \ 2l \ 2\bar{l} \rightarrow 2d \ 2\bar{d} \ 4l \ 4\bar{l}$ will quickly thermalize and not have any impact on the Helium production. Requiring $\tau_\chi \leq 10^{-2} \text{ sec}$., we get, $V_{BL} \leq 10^{11} \text{GeV}$ for $\delta \simeq 0.1$. Thus, in the presence of CP-violating Planck effects, the results of Ref.[4] are extended as follows:

$$V_{BL} \leq \left(\frac{m_\nu}{25\text{eV}} \right)^{4/7} \times 10 \text{ TeV}, \quad (7.a)$$

and

$$10^8 \text{GeV} \leq V_{BL} \leq 10^{11} \text{GeV}. \quad (7.b)$$

A few comments are in order: there can also be heavy decay in the CP-conserving limit such as $\chi \rightarrow t\bar{t}$, $b\bar{b}$ etc. Their effect on these bounds is negligible.

Another point of interest is that the decay of the massive Majoron will generate large amount of entropy if $\tau_\chi \geq 10^{-12}$ sec. (corresponding to $m_\chi \geq 10^3 \text{GeV}$). This entropy generation will dilute any preexisting baryon asymmetry of the universe by a factor x , which can be obtained from energy conservation. For $m_\chi \gg T_\chi$, we find

$$x \equiv \frac{S_f}{S_i} \simeq \left(\frac{120Y}{\pi^4 g^*} \frac{m_\chi}{T_\chi} \right)^{3/4}, \quad (8)$$

where $T_\chi \simeq (1/\tau_\chi \text{ in sec.})^{1/2} \text{MeV}$, and g^* is the number of degrees of freedom at $T = T_\chi$. For instance, for $m_\chi \simeq 10^3 \text{TeV}$, $\tau_\chi \simeq 10^{-5} \text{sec.}$ leads to $x \simeq 10^3$. This implies that any baryon or lepton asymmetry generated earlier than this epoch must be of order $10^{-6} n_\gamma$ in order to explain present observation. Otherwise, the observed asymmetry must be generated at a later time *i.e.* $t_U > \tau_\chi$.

In conclusion, if the non-perturbative Planck scale induced operators are CP-violating, there appears an allowed high mass scale window for the scale of global B-L symmetry breaking (*i.e.* $10^8 \text{GeV} \leq V_{BL} \leq 10^{11} \text{GeV}$) in addition to the allowed range below 10 TeV obtained in Ref.[4].

References

- [1] R. Holman, S. Hsu, T. Kephart, E. Kolb, R. Watkins and L. Widrow, Phys. Lett. B282 (1992) 132; M. Kamionkowski and J. March-Russel, Phys. Lett. B282 (1992) 137; S. Barr and D. Seckel, Phys. Rev. D46 (1992) 538.
- [2] see, C. Misner, K. Thorne and J. Wheeler, *Gravitation* (Freeman, San Francisco, 1973).
- [3] Y. Chikashige, R.N. Mohapatra and R.D. Peccei, Phys. Lett. B98 (1980) 265.
- [4] E. Akhmedov, Z. Berezhiani, R. Mohapatra and G. Senjanović , Phys. Lett. B299 (1993) 90.
- [5] J. Cline, K. Kainulainen and K. Olive, UMN-TH-1113-93, TPI-MINN-93/13-T, April 1993.
- [6] G. Steigman, K. Olive and D. Schramm, Phys. Rev. Lett. 43 (1979) 239; K. Olive, D. Schramm and G. Steigman, Nucl. Phys. B180 (1981) 497.